

CS 2336: Discrete Mathematics


Instructor: Cheng-Hsin Hsu

Course Information

- **TextBook:** Discrete and Combinatorial Mathematics, R. Grimaldi, 5th Ed., Addison Wesley
- **Lecture:** Mondays 3:30 p.m.-5:20 p.m. and Thursdays 2:20-3:10 p.m.
- **Location:** 台達館 103
- **Office Hour:** Thursdays 3:10 p.m. – 4:00 p.m.
- **Course Website:**
<http://nmsl.cs.nthu.edu.tw/index.php/courses>
- **TA:** Tao-Ya Fan-Jiang (tyfanchiang92@gmail.com)
- **TA Office Hour:** Mondays 5:30 – 6:30 p.m. at Delta 713

Topics We Plan to Cover

1. **Fundamental Principles of Counting (Chapter 1 in textbook)**
2. **Fundamental of Logic (Chapter 2)**
3. **Set Theory (Chapter 3)**
4. **Properties of Integers: Mathematical Induction (Chapter 4)**
5. **Relations and Functions (Chapter 5)**
6. **Languages: Finite State Machines (Chapter 6)**
7. **Relations: The Second Time Around (Chapter 7)**
8. **The Principle of Inclusion and Exclusion (Chapter 8)**
9. **Generating Functions (Chapter 9)**
10. **Recurrence Relations (Chapter 10)**
11. **An Introduction to Graph Theory (Chapter 11)**
12. **Trees (Chapter 12)**
13. **Optimization and Matching (Chapter 13)**



Not Covered: Applied Algebra

Grading Policy

- **Quizzes (60%):** Ten times, One for each chapter
 - Given on Mondays, either at the beginning or end of the lecture
 - Sample questions will be given as homework, which are not collected nor graded
 - No makeup quizzes, unless an email requesting for a leave is sent to and approved by the instructor **before** each quiz
- **Midterm Exam (20%):** 2-hr exam on Apr. 20th
- **Final Exam (20%):** 2-hr exam on Jun. 22th

Tentative Schedule

Week	Mondays	Thursdays	Homework/Quiz Solutions
1: Feb 22		Introduction	
2: Mar 1	Ch. 1 Fundamental Principles of Counting	Ch. 2 Fundamental of Logic	
3: Mar 8	Ch. 2 Fundamental of Logic	Ch. 2 Fundamental of Logic	
4: Mar 15	Ch. 2 Fundamental of Logic	Conference Travel	
5: Mar 22	Conference Travel	Ch. 4 Properties of Integers: Mathematical Induction	
6: Mar 29	Ch. 4 Properties of Integers: Mathematical Induction	Ch. 5 Relations and Functions	
7: Apr 5	Holiday	Ch. 5 Relations and Functions	
8: Apr 12	Ch. 5 Relations and Functions	Ch. 6 Finite State Machines	
9: Apr 19	Mid Term Exam (Chs. 1 - 5)	Conference Travel (Mid Term Review)	
10: Apr 26	Ch. 7 Relations: The Second Time Around	Ch. 7 Relations: The Second Time Around	
11: May 3	Ch. 7 Relations: The Second Time Around	Ch. 7 Relations: The Second Time Around	
12: May 10	Ch. 8 The Principle of Inclusion and Exclusion	Ch. 8 The Principle of Inclusion and Exclusion	
13: May 17	Ch. 8 The Principle of Inclusion and Exclusion	Ch. 8 The Principle of Inclusion and Exclusion	
14: May 24	Ch. 9 Generating Functions	Ch. 9 Generating Functions	
15: May 31	Ch. 9 Generating Functions	Ch. 10 Recurrence Relations	
16: Jun 7	Ch. 10 Recurrence Relations	Ch. 10 Recurrence Relations	
17: Jun 14	Ch. 11 An Introduction to Graph Theory	Ch. 11 An Introduction to Graph Theory	
18: Jun 21	Final Exam (Chs. 6 - 11)		

What is Discrete Mathematics?

- Covers various kinds of topics
 - Logics
 - Combinatorial
 - Algorithms
 - Graph Theory
 - Number Theory
- Discrete ← something you can count

What is a Proof?

- Vaguely speaking:
 - To convince someone that something is true
- Mathematical proof:
 - To show if some axioms are true, then some statements are also true
- What we will do in this course is somewhere between these two definitions
 - You need to get the main idea
 - Details are not important

Examples of Proofs (1/2)

- Prove that there are infinite prime numbers [Eulclid 300 BC]
 - A prime is a nature number with exactly two divisors
 - Assume there are finite # of primes: p_1, p_2, \dots, p_n , sorted in the increasing order
 - We create a new prime $p = p_1 p_2 \cdots p_n + 1$
 - Why p is a prime? $\rightarrow p / p_i$ for any i has remainder 1
 - Moreover, p is larger than p_1, p_2, \dots, p_n , so p is a new prime \rightarrow contradicting to the assumption that there are only n prime numbers out there
 - Hence, there are infinite number of prime numbers

Examples of Proofs (2/2)

- Prove that $\sqrt{2}$ is irrational (by contradiction)
 - Assume $\sqrt{2}$ is a rational number, then we write $\sqrt{2} = p/q$, where p and q are the lowest terms
 - We know $2 = p^2/q^2$ and $2q^2 = p^2 \rightarrow p$ is even
 - We write $p=2k$, then we have $2q^2 = 4k^2$ and $q^2 = 2k^2$
 - Then, q is also even \rightarrow contradiction!
 - Hence we know $\sqrt{2}$ is irrational

Questions?

Questions so far?

CS 2336: Discrete Mathematics

Chapter 1

Fundamental Principles of Counting

Instructor: Cheng-Hsin Hsu

Rule of Sum

- Q: There are 6, 8, and 12 introductory books for C++, Java, and Perl, respectively. If Joe wants to learn a first language, how many choices (of books) does he have?
- Q: Say Joe has two friends, who own 3 and 5 of these books. How many unique books can Joe borrow from his friends?
- Rule: If task 1 can be performed in m ways, task 2 can be performed in n ways, and the two tasks cannot be performed simultaneously, then there are $m+n$ ways to perform either task 1 or 2.

Rule of Product

- Q: There is a singer try-out with 4 men and 6 women, how many different way to form a couple of singers?
- Q: How many different license plates of two letters followed by four digits can we produce with/without repetition
- Rule: If task 1 can be performed in m ways, task 2 can be performed in n ways, there are mn ways to sequentially perform tasks 1 and 2.

Combinations of Two Rules

- Q: A cafeteria offers 6 kinds of muffins, 8 kinds of sandwiches, and 3 different beverages, and each combo consists of a muffin (or sandwiches) and a beverage. How many different combos does this cafeteria offer?

Permutation

- Q: From a class of 10 students, choose and seat 5 of them in a line for a picture. **How many linear arrangement are possible?**
 - Write the solution using n factorial
- Definition: Given a collection of n distinct objects, any arrangement of these objects is called a permutation of the collection
- For n distinct objects and an integer $1 \leq r \leq n$, per the rule of product, the number of permutation of size r of n object is:
$$P(n, r) = \frac{n!}{(n - r)!}$$

Repeated Permutation

- Q: How many 5-character permutations of the letters in “COMPUTER” if
 - Repetitions are not allowed
 - Repetitions are allowed
- Q: How many permutations of the letters in
 - BALL
 - DATABASES
- General Result: If there are n objects with n_1 (indistinguishable) type 1 objects, ..., n_r type r objects, there are $\frac{n!}{n_1!n_2!\cdots n_r!}$ arrangements of the given n objects

Nonlinear Permutation

- Q: If six students are seated at a round table, how many different arrangements are possible?
 - Assume that arrangements are considered the same when one can be derived by rotating the other
- Q: How many ways we can arrange 3 males and 3 female around a table so that the sexes alternate?

Combinations

- Q: How many possible permutations of 3 from 5 cards? What if the cards are considered as **unordered**?

- General rule: For n distinct object, the number of combinations of r object, where $0 \leq r \leq n$, is:

$$C(n, r) = \frac{P(n, r)}{r!} = \frac{n!}{r!(n - r)!}$$

- Q: How many different ways a student can answer 7 out of 10 questions in a test?
- Q: What if the student needs to pick 3 questions from the first 5 questions and 4 questions from the last 5 questions?

Permutation and/or Combination

- Q: How many arrangements of the letters in TALLAHASSEE with no adjacent A's?
 - Requires both permutation and combination
- Q: The coach wants to form four teams of 9 students each from a class of 36 students. Call the teams A, B, C, and D, how many different ways can the coach form the teams?
 - Solve it using combinations
 - Solve it using permutations

Summation

- We use Greek symbol sigma to represent summations

- $\sum_{i=m}^n a_i = a_m + a_{m+1} + \cdots + a_n$

- i is the index

- m is the lower limit

- n is the upper limit

- Example:

$$\sum_{i=2}^3 \binom{5}{7-j} \binom{5}{j} = \binom{5}{5} \binom{5}{2} + \binom{5}{4} \binom{5}{3}$$

Another Example

- Q: Let n be a positive integer, there are 3^n strings of an alphabet consisting of symbols 0, 1, and 2. Define $\text{weight}(x) = x_1 + x_2 + \dots + x_n$, for $x = x_1 x_2 \dots x_n$. For $n=10$, how many strings with even weights?

Theorems

- Lemma: Prove that for two integers n and r , $n \geq r \geq 0$, we have
$$\binom{n}{r} = \binom{n}{n-r}$$

- Theorem (Binomial): Prove that

$$(x + y)^n = \binom{n}{0} x^0 y^n + \binom{n}{1} x^1 y^{n-1} + \dots + \binom{n}{n} x^n y^0 = \sum_{k=0}^n \binom{n}{k} x^k y^{n-k}$$

- $\binom{n}{k}$ is called binomial coefficient

- Corollary

- $2^n = \binom{n}{0} + \binom{n}{1} + \dots + \binom{n}{n}$
- $0 = \binom{n}{0} - \binom{n}{1} + \dots + (-1)^n \binom{n}{n}$

Theorems (cont.)

- Theorem (Multinomial): Prove that the coefficient of $x_1^{n_1} x_2^{n_2} \dots x_t^{n_t}$ in the expansion of $(x_1 + x_2 + \dots + x_t)^n$ is
$$\frac{n!}{n_1! n_2! \dots n_t!}$$
- $\binom{n}{n_1, n_2, \dots, n_t}$ is called multinomial coefficient
- Example: Determine the coefficient of $a^2 b^3 c^2 d^5$ in the expansion of $(a + 2b - 3c + 2d + 5)^{16}$

Comb. with Repetition

- Q: Seven students stop at a fast food restaurant where each of them can order a burger, a hot dog, a taco, or a sandwich. The restaurant only cares about how many burgers, hot dogs, tacos, and sandwiches do the students order. What is the number of possible solutions?
- General Rule: The number of combinations of, with repetition, r objects from n distinct objects is:

$$\frac{(n + r - 1)!}{r!(n - 1)!} = \binom{n + r - 1}{r}$$

Examples

- Q: A donut shop offer 20 kinds of donuts. Assuming that there are plenty of donuts of each kind, how many ways for a kid to buy a dozen donuts?
- Q: How many solutions does $x_1 + x_2 + x_3 + x_4 = 7$ have, for positive $x_1, x_2, x_3, \text{ and } x_4$?
- Q: A father distributes \$1000 among 4 kids at a step of \$100
 - How many different ways to distribute \$1000 if some kids may get nothing?
 - How many ways to distribute \$1000 if each kid is guaranteed to have at least \$100?

Equivalence

- The number of selections, with repetition, of size r from a collection of size n
- The number of ways r identical objects can be distributed among n distinct containers
- The number of integer solutions of the equation

$$x_1 + x_2 + \cdots + x_n = r, \quad x_i \geq 0, 1 \leq i \leq n$$

Examples

- Q: How many nonnegative solutions does the inequality $x_1 + x_2 + \cdots + x_6 < 10$ have?
- Q: How many times the print statement is executed?
for i = 1 to 20
 for j = 1 to i
 for k = 1 to j
 print (i+j+k);

Take-home Exercise

- Exercise 1.1 and 1.2: 15, 22, 28, 32, 33
- Exercise 1.3: 13, 16, 25, 29, 34
- Exercise 1.4: 7, 17, 24, 26, 28