

SOLUTION

Ex 7.1: 1, 5, 6, 9, 17

Ex 7.2: 4, 14, 17, 18, 26

Ex 7.3: 1, 7, 18, 23, 25

Ex 7.4: 2, 6, 7, 12, 14

Ex 7.5: 1, 3

Ex 7.1: (1)

- a) $\{(1,1), (2,2), (3,3), (4,4), (1,2), (2,1), (2,3), (3,2)\}$
- b) $\{(1,1), (2,2), (3,3), (4,4), (1,2)\}$
- c) $\{(1,1), (2,2), (1,2), (2,1)\}$

Ex 7.1: (5)

- a) reflexive, antisymmetric, transitive
- b) transitive
- c) reflexive, symmetric, transitive
- d) symmetric
- e) (odd): symmetric
- f) (even): reflexive, symmetric, transitive
- g) reflexive, symmetric
- h) reflexive, transitive

Ex 7.1: (6)

- The relation in part (a) is a partial order.
The relation in parts (c) and (f) are equivalence relations.

Ex 7.1: (9)

- a) False: Let $A = \{1,2\}$ and $\mathcal{R} = \{(1,2), (2,1)\}$.
- b)
 - (i) Reflexive: True.
 - (ii) Symmetric: False.
Let $A = \{1,2\}$, $\mathcal{R}_1 = \{(1,1)\}$, $\mathcal{R}_2 = \{(1,1), (1,2)\}$
 - (iii) Antisymmetric and transitive: False.
Let $A = \{1,2\}$, $\mathcal{R}_1 = \{(1,2)\}$, $\mathcal{R}_2 = \{(1,2), (2,1)\}$
- c)
 - (i) Reflexive: False.
Let $A = \{1,2\}$, $\mathcal{R}_1 = \{(1,1)\}$, $\mathcal{R}_2 = \{(1,1), (2,2)\}$
 - (ii) Symmetric: False.
Let $A = \{1,2\}$, $\mathcal{R}_1 = \{(1,2)\}$, $\mathcal{R}_2 = \{(1,2), (2,1)\}$
 - (iii) Antisymmetric: True.
 - (iv) Transitive: False.
Let $A = \{1,2\}$, $\mathcal{R}_1 = \{(1,2), (2,1)\}$,
 $\mathcal{R}_2 = \{(1,1), (1,2), (2,1), (2,2)\}$
- d) True.

Ex 7.1: (17)

a) $\binom{7}{5} \binom{21}{0} + \binom{7}{3} \binom{21}{1} + \binom{7}{0} \binom{21}{2}$

b) $\binom{7}{5} \binom{21}{0} + \binom{7}{3} \binom{21}{1} + \binom{7}{1} \binom{21}{2}$

Ex 7.2: (4)

- a) $\mathcal{R}_1 \circ (\mathcal{R}_2 \cup \mathcal{R}_3) = \{(1,4), (1,5), (3,4), (3,5), (2,6), (1,6)\}$
 $(\mathcal{R}_1 \circ \mathcal{R}_2) \cup (\mathcal{R}_1 \circ \mathcal{R}_3) = \{(1,4), (1,5), (1,6), (2,6), (3,4), (3,5)\}$
- b) $\mathcal{R}_1 \circ (\mathcal{R}_2 \cup \mathcal{R}_3) = \{(1,5), (3,5)\}$
 $(\mathcal{R}_1 \circ \mathcal{R}_2) \cup (\mathcal{R}_1 \circ \mathcal{R}_3) = \{(1,4), (1,5), (3,5)\}$

Ex 7.2: (14.1)

```
10! THIS PROGRAM MAY BE USED TO DETERMINE IF A RELATION
20! ON A SET OF SIZE N, WHERE N <= 20, IS AN
30! EQUIVALENCE RELATION. WE ASSUME WITHOUT LOSS OF
40! GENERALITY THAT THE ELEMENTS ARE 1,2,3,...,N.
50!
60 INPUT "N ="; N
70 PRINT " INPUT THE RELATION MATRIX FOR THE RELATION"
80 PRINT "BEING EXAMINED BY TYPING A(I,J) = 1 FOR EACH"
90 PRINT "1 <= I <= N, 1 <= J <= N, WHERE (I,J) IS IN"
100 PRINT "THE RELATION. WHEN ALL THE ORDERED PAIRS HAVE"
110 PRINT "BEEN ENTERED TYPE 'CONT' "
120 STOP
130 DIM A(20,20), C(20,20), D(20,20)
140 FOR K = 1 TO N
150     T = T + A(K,K)
160 NEXT K
170 IF T = N THEN &
180     PRINT "R IS REFLEXIVE"; X = 1: GO TO 190
PRINT "R IS NOT REFLEXIVE"
```

Ex 7.2: (14.2)

```
190 FOR I = 1 TO N
200     FOR J = I + 1 TO N
210         IF A(I,J) <> A(J,I) THEN GO TO 260
220     NEXT J
230 NEXT I
240 PRINT "R IS SYMMETRIC"; Y = 1
250 GO TO 270
260 PRINT "R IS NOT SYMMETRIC"
270 MAT C = A
280 MAT D = A*C
290 FOR I = 1 TO N
300     FOR J = 1 TO N
310         IF D(I,J) > 0 AND A(I,J) = 0 THEN GO TO 360
320     NEXT J
330 NEXT I
340 PRINT "R IS TRANSITIVE"; Z = 1
350 GO TO 370
360 PRINT "R IS NOT TRANSITIVE"
370 IF X + Y + Z = 3 THEN &
            PRINT "R IS AN EQUIVALENCE RELATION" &
            ELSE PRINT "R IS NOT AN EQUIVALENCE RELATION"
380 END
```

Ex 7.2: (17.i)

- $\mathcal{R} = \{(a, b), (b, a), (a, e), (e, a), (b, c), (c, b), (b, d), (d, b), (b, e), (e, b), (d, e), (e, d), (d, f), (f, d)\}$

- $M(\mathcal{R}) = \begin{bmatrix} 0 & 1 & 0 & 0 & 1 & 0 \\ 1 & 0 & 1 & 1 & 1 & 0 \\ 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 1 & 1 \\ 1 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \end{bmatrix}$

Ex 7.2: (17.ii)

- $\mathcal{R} = \{(a, b), (b, e), (d, b), (d, c), (e, f)\}$

- $M(\mathcal{R}) = \begin{bmatrix} 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$

Ex 7.2: (17.iii)

- $\mathcal{R} = \{(a, a), (a, b), (b, a), (c, d), (d, c), (d, e), (e, d), (d, f), (f, d), (e, f), (f, e)\}$

- $M(\mathcal{R}) = \begin{bmatrix} 1 & 1 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 1 & 1 \\ 0 & 0 & 0 & 1 & 0 & 1 \\ 0 & 0 & 0 & 1 & 1 & 0 \end{bmatrix}$


Ex 7.2: (17.iv)

- $\mathcal{R} = \{(b, a), (b, c), (c, b), (b, e), (c, d), (e, d)\}$

- $M(\mathcal{R}) = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 1 & 0 & 1 & 0 \\ 0 & 1 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}$


Ex 7.2: (18.a)

- $\mathcal{R} = \{(v, w), (v, x), (w, v), (w, x), (w, y), (w, z), (x, z), (y, z)\}$



Ex 7.2: (18.b)

- $\mathcal{R} = \{(v, w), (v, x), (v, y), (w, v), (w, x), (x, v), (x, w), (x, z), (y, v), (y, z), (z, x), (z, y)\}$



Ex 7.2: (26.a, 26.b)

a) Let $k \in \mathbb{Z}^+$. Then


$R^{12k} = \{(1,1), (2,2), (3,3), (4,4), (5,5), (6,6), (7,7)\}$ and
 $R^{12k+1} = R$. The smallest value of $n > 1$ such that
 $R^n = R$ is $n = 13$. For all multiples of 12 the graph
consists of all loops. When $n = 3$, $(5,5)(6,6)(7,7) \in R^3$,
and this is the smallest power of R that contains at least
one loop.

b) When $n = 2$, we find $(1,1), (2,2)$ in R . For all $k \in \mathbb{Z}^+$,
 $R^{30k} = \{(x,x) | x \in \mathbb{Z}^+, 1 \leq x \leq 10\}$ and $R^{30k+1} = R$.
Hence R^{31} is the smallest power of R (for $n > 1$) where
 $R^n = R$.

Ex 7.2: (26.c)


- Let R be a relation on set A where $|A| = m$. Let G be the directed graph associated with R – each component of G is a directed cycle C_i on m_i vertices, with $1 \leq i \leq k$. (Thus $m_1 + m_2 + \dots + m_k = m$.) The smallest power of R where loops appear is R^t , for $t = \min\{m_i | 1 \leq i \leq k\}$.
- Let $s = \text{lcm}(m_1, m_2, \dots, m_k)$. Then R^{rs} =the identity (equality) relation on A and $R^{rs+1} = R$, for all $r \in \mathbb{Z}^+$. The smallest power of R that reproduces R is $s + 1$.

Ex 7.3: (1)



Ex 7.3: (7)

a)



- b) $3 < 2 < 1 < 4$ or $3 < 1 < 2 < 4$
- c) 2

Ex 7.3: (18)

- a) (i) Only one such upper bound – $\{1,2,3\}$.
(ii) Here the upper bound has the form $\{1,2,3, x\}$ where $x \in \mathcal{U}$ and $4 \leq x \leq 7$. Hence there are four such upper bounds.
(iii) There are $\binom{4}{2}$ upper bounds of B that contain five elements from \mathcal{U} .
- b) $\binom{4}{0} + \binom{4}{1} + \binom{4}{2} + \binom{4}{3} + \binom{4}{4} = 2^4 = 16$
- c) $lub B = \{1,2,3\}$
- d) One—namely \emptyset
- e) $glb B = \emptyset$

Ex 7.3: (23)

- a) Flase. Let $U = \{1,2\}$, $A = P(U)$, and \mathcal{R} be the inclusion relation. Then (A, \mathcal{R}) is a lattice where for all $S, T \in A$, $\text{lub}\{S, T\} = S \cup T$ and $\text{glb}\{S, T\} = S \cap T$. However, $\{1\}$ and $\{2\}$ are not related, so (A, \mathcal{R}) is not a total order.
- b) If (A, \mathcal{R}) is a total order, then for all $x, y \in A$, $x\mathcal{R}y$ or $y\mathcal{R}x$. For $x\mathcal{R}y$, $\text{lub}\{x, y\} = y$ and $\text{glb}\{x, y\} = x$. Consequently, (A, \mathcal{R}) is a lattice.

Ex 7.3: (25)

- a) a
- b) a
- c) c
- d) e
- e) z
- f) e
- g) v

(A, \mathcal{R}) is a lattice with z the greatest (and only maximal) element and a the least (and only minimal) element.

Ex 7.4: (2)

- a) There are three choices for placing 8 – in either A_1 , A_2 or A_3 . Hence there are three partitions of A for the conditions given.
- b) There are two possibilities with $7 \in A_1$, and two others with $8 \in A_1$. Hence there are four partitions of A under these conditions.
- c) If we place 7,8 in the same cell for a partition we obtain three of the possibilities. If not, there are three choices of cells for 7 and two choices of cells for 8 – and six more partitions that satisfy the stated restrictions. In total – by the rules of sum and produce – there are $3 + (3)(2) = 3 + 6 = 9$ such partitions.

Ex 7.4: (6)

- a) For all $(x, y) \in A$, since $x = x$, it follows that $(x, y)R(x, y)$, so R is *reflexive*. If $(x_1, y_1), (x_2, y_2) \in A$ and $(x_1, y_1)R(x_2, y_2)$, then $x_1 = x_2$, so $x_2 = x_1$ and $(x_2, y_2)R(x_1, y_1)$. Hence R is *symmetric*. Finally, let $(x_1, y_1), (x_2, y_2), (x_3, y_3) \in A$ with $(x_1, y_1)R(x_2, y_2)$ and $(x_2, y_2)R(x_3, y_3)$. $(x_1, y_1)R(x_2, y_2) \Rightarrow x_1 = x_2$; $(x_2, y_2)R(x_3, y_3) \Rightarrow x_2 = x_3$. With $x_1 = x_2, x_2 = x_3$, it follows that $x_1 = x_3$, so $(x_1, y_1)R(x_3, y_3)$ and R is *transitive*.
- b) Each equivalence class consists of the points on a vertical line. The collection of these vertical lines then provides a partition of the real plane.

Ex 7.4: (7)

- a) For all $(x, y) \in A$, $x + y = y + x \Rightarrow (x, y)R(x, y)$.
- $(x_1, y_1)R(x_2, y_2) \Rightarrow x_1 + y_1 = x_2 + y_2 \Rightarrow x_2 + y_2 = x_1 + y_1$
 $\Rightarrow (x_2, y_2)R(x_1, y_1)$. $(x_1, y_1)R(x_2, y_2), (x_2, y_2)R(x_3, y_3)$
 $\Rightarrow x_1 + y_1 = x_2 + y_2, x_2 + y_2 = x_3 + y_3$, so $x_1 + y_1 = x_3 + y_3$
and $(x_1, y_1)R(x_3, y_3)$.
- Since R is reflexive, symmetric and transitive, it is an equivalence relation.
- b) $[(1,3)] = \{(1,3), (2,2), (3,1)\};$
 $[(2,4)] = \{(1,5), (2,4), (3,3), (4,2), (5,1)\};$ $[(1,1)] = \{(1,1)\}.$
- c) $A = \{(1,1)\} \cup \{(1,2), (2,1)\} \cup \{(1,3), (2,2), (3,1)\} \cup$
 $\{(1,4), (2,3), (3,2), (4,1)\} \cup \{(1,5), (2,4), (3,3), (4,2), (5,1)\} \cup$
 $\{(2,5), (3,4), (4,3), (5,2)\} \cup \{(3,5), (4,4), (5,3)\} \cup \{(4,5), (5,4)\} \cup$
 $\{(5,5)\}.$

Ex 7.4: (12)

a) $2^{10} = 1024$

b) $\sum_{i=1}^5 S(5, i) = 1 + 15 + 25 + 10 + 1 = 52$

c) $1024 - 52 = 972$

d) $S(5, 2) = 15$

e) $\sum_{i=1}^4 S(4, i) = 1 + 7 + 6 + 1 = 15$

f) $\sum_{i=1}^3 S(3, i) = 1 + 3 + 1 = 5$

g) $\sum_{i=1}^3 S(3, i) = 1 + 3 + 1 = 5$

h) $(\sum_{i=1}^3 S(3, i)) - (\sum_{i=1}^2 S(2, i)) = 3$

Ex 7.4: (14)

- a) Not possible. With R reflexive, $|R| \leq 7$.
- b) $R = \{(x, x) | x \in \mathbb{Z}, 1 \leq x \leq 7\}$.
- c) Not possible. With R symmetric, $|R| - 7$ must be even.
- d) $R = \{(x, x) | x \in \mathbb{Z}, 1 \leq x \leq 7\} \cup \{(1,2), (2,1)\}$.
- e) $R = \{(x, x) | x \in \mathbb{Z}, 1 \leq x \leq 7\}$
 $\quad \cup \{(1,2), (2,1)\} \cup \{(3,4), (4,3)\}$.
- f) Not possible with $r - 7$ odd.
- g) Not possible. See the remark at the end of Section 7.4.
- h) Not possible with $r - 7$ odd.
- i) Not possible. See the remark at the end of Section 7.4.

Ex 7.5: (1)

a) $P_1: \{s_1, s_4\}, \{s_2, s_3, s_5\}$

$(v(s_1, 0) = s_4)E_1(v(s_4, 0) = s_1)$ but

$(v(s_1, 1) = s_1)E_1(v(s_4, 1) = s_3)$, so $s_1 \not E_2 s_4$.

$(v(s_2, 1) = s_3)E_1(v(s_3, 1) = s_4)$, so $s_2 \not E_2 s_3$.

$(v(s_2, 0) = s_3)E_1(v(s_5, 1) = s_3)$ and

$(v(s_2, 1) = s_3)E_1(v(s_5, 1) = s_3)$, so $s_2 \not E_2 s_5$.

Since $s_2 \not E_2 s_3$ and $s_2 \not E_2 s_5$, it follows that $s_3 \not E_2 s_5$.

Hence P_2 is given by $P_2: \{s_1\}, \{s_2, s_5\}, \{s_3\}, \{s_4\}$.

$(v(s_2, x) = s_3)E_2(v(s_5, x) = s_3)$ for $x = 0, 1$.

Hence $s_2 E_3 s_5$ and $P_2 = P_3$.

Consequently, states s_2 and s_5 are equivalent.

b) States s_2 and s_5 are equivalent.

c) States s_2 and s_7 are equivalent; s_3 and s_4 are equivalent

Ex 7.5: (3)

- a) s_1 and s_7 are equivalent; s_4 and s_5 are equivalent
- b) (i) 0000
(ii) 0
(iii) 00

M:	v		w	
	0	1	0	1
s_1	s_4	s_1	1	0
s_2	s_1	s_2	1	0
s_3	s_6	s_1	1	0
s_4	s_3	s_4	0	0
s_6	s_2	s_1	1	0